arXiv:gr-qc/9812026v2 30 Dec 1998

Static Cylindrically Symmetric Solution

Mihaela Time

6th December 1998

Abstract

Here is described a static axisymmetric solution with an additional cylindrical symmetry for which the matter consists in a cosmological and a dust term.

1 The field equations

Stationary gravitational fields are characterized by the existence of a timelike Killing vector field ξ. So, in a stationary space-time $(M,\, g)$ we can have a global causal structure, i.e. we can introduce an adapted coordinate system $(x^a) = (x^\alpha,\, t)$, $\xi = \frac{\partial}{\partial t}$ so that the metric g_{ab} is independent of t,

$$(1) \qquad \mathrm{d}s^2 = h_{\alpha\beta}\mathrm{d}x^\alpha \mathrm{d}x^\beta + F(\mathrm{d}t + A_\alpha \mathrm{d}x^\alpha)^2, \qquad F \equiv \xi_a\xi^a < 0.$$

The timelike unitary vector field $h^0 \equiv (-F)^{-\frac{1}{2}}\xi$, which is globally defined on M, indicates not only the time-orientation in every point $p \in M$, but also gives a global time coordinate t on M.

Stationarity (i.e. time translation symmetry) means that there exists a 1-dimensional group G_1 of isometries ϕ_t whose orbits are timelike curves parametrized by t. The smooth surjective submersion

$$\pi' : M \longrightarrow \mathcal{S}_3, \quad p \in M \longmapsto \pi'(p) = \phi_t(p),\ \ \phi_0(p) = p,\ \ \dot{\phi}_p(t) = \xi(\phi_p(t))$$

over the 3-dimensional differentiable factor manifold $\mathcal{S}_3$, whose elements are the orbits of G_1 determines a fibration on M; the metric g_{ab} is constant on the fibres of π' (integral curves of ξ) and there is a 1:1 correspondence between tensors $T = (T^{a\ldots}_{b\ldots})$on M satisfying

$$\xi^a T^{b\ldots}_{a\ldots} = \xi_b T^{b\ldots}_{a\ldots} = 0 \text{ and } \pounds_\xi T = 0$$

and the tensors on $\mathcal{S}_3$. So using the 3-projection formalisme (by Geroch(1971)) developed on $\mathcal{S}_3$ (see [3]), the Einstein's field equations

$$(2) \qquad R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = \kappa T_{ab},$$

will take the following simplified form for stationary fields:

$$(3)\begin{cases} R^{(3)}_{ab} = \frac{1}{2} F^{-2} (\frac{\partial F}{\partial x^a} \frac{\partial F}{\partial x^b} + \omega_a \omega_b) + \kappa (h^c_a h^d_b - F^{-2} \tilde{h}_{ab} \xi^a \xi^b)(T_{cd} - \frac{1}{2} T g_{cd}); \\ F^{\|a}_{,a} = F^{-1} \tilde{h}_{ab} (\frac{\partial F}{\partial x^a} \frac{\partial F}{\partial x^b} - \omega_a \omega_b) - 2\kappa F^{-1} \xi^a \xi^b (T_{ab} - \frac{1}{2} T g_{ab}); \\ \omega^{\|a}_a = 2F^{-1} \tilde{h}_{ab} \frac{\partial F}{\partial x^a} \omega_b \\ F \epsilon^{abc} \omega_{c,b} = 2\kappa h^a_b T^b_c \xi^c \end{cases}$$

Here, "$\|$" denotes the covariant derivative associated with the conformal metric tensor $\tilde{h}_{ab} = -F h_{ab}$ on $\mathcal{S}_3$ ($h_{ab} = g_{ab} + h^0_a h^0_b$ is the projection tensor), $\epsilon^{abc} = \epsilon^{dabc} h^0_d$ and $\omega^a = \frac{1}{2} \epsilon^{abcd} \xi_b \xi_{c;d} \neq 0$ is the rotational vector ($\omega^a \xi_a = 0$, $\pounds_\xi \omega = 0$).

The circularity theorem (due to Kundt) states that an (axisymmetric) metric can be written [1] in the (2+2)-split if and only if the conditions

$$(4) \qquad (\eta^{[a} \xi^b \xi^{c;d]})_{;e} = 0 = (\xi^{[a} \eta^b \eta^{c;d]})_{;e}$$

are satisfied [1].

The existence of the orthogonal 2-surfaces is possible for dust solutions, provided that the 4-velocity of dust satisfies the condition:

$$(5) \qquad u_{[a} \xi_b \eta_{c]} = 0, \quad u^a = (-H)^{-\frac{1}{2}} (\xi^a + \Omega \eta^a) = (-H)^{-\frac{1}{2}} l^i \xi^a_i, \quad \text{where}$$
$$l^i \equiv (1, \Omega), \quad H = \gamma_{ij} l^i l^j, \quad \gamma_{ij} \equiv \xi^a_i \xi_{aj}, \quad i,j = 1,2, \quad \xi_1 = \xi; \ \xi_2 = \eta$$

i.e., the trajectories of the dust lie on the transitivity surfaces of the group generated by the Killing vectors ξ, η. Here Ω is the angular velocity of the matter with respect to infinity.

Using an adapted coordinate system, the metric (1) can be written in the following form:

$$(6) \qquad \mathrm{d}s^2 = e^{-2U} [e^{2V} (\mathrm{d}r^2 + \mathrm{d}z^2) + W^2 \mathrm{d}\varphi^2] - e^{2U} (\mathrm{d}t + A \mathrm{d}\varphi)^2$$

---

[1] Here I use the convention: round brackets denote symmetrization and square brackets antisymmetrization.

where the functions [2] U, V, W and A depend only on the coordinates (r, z); these coordinates are also conformal flat coordinates on the 2-surface S_2 orthogonal to 2-surface T_2 of the commuting Killing vectors $\xi = \partial_t$ and $\eta = \partial_\varphi$.

If we identify the 4-velocity of the dust u^a with timelike Killing vector $\xi^a = \partial_t = (0,0,0,1)$ then (6) represents a co-moving system $(x^1 = r,\ x^2 = z,\ x^3 = \varphi,\ x^0 = t)$ with dust, $u_a = \xi_a = (0,\ 0,\ -e^{2U}A,\ -e^{2U})$ and

$$\left\{\begin{array}{l} g_{11} = g_{22} = e^{-2U+2V} = h_{11} = h_{22}, \\ g_{33} = e^{-2U}W^2 - e^{2V}A^2 = h_{33}, \quad g_{00} = \xi_0 = -e^{2U} = F, \\ g_{03} = \xi_3 = -e^{2U}A, \quad g_{13} = g_{23} = g_{10} = g_{20} = 0 \end{array}\right. \tag{7}$$

The energy-momentum tensor T_{ab} has the form as:

$$\kappa T_{ab} = -\Lambda g_{ab} + \mu u_a u_b, \quad \mu > 0, \Lambda = \text{const.} > 0. \tag{8}$$

The conservation law, $T^{ab}_{;b} = 0$ implies $U_{,a} = 0$ and is a consequence of the field equations, being used in place of one of the equations (3).

If $\Omega = \text{const.}$, i.e., for rigid rotation we have

$$\sigma = 0 = \theta \Leftrightarrow u_{(a;b)} + u_{(a}\dot{u}_{b)} = 0 \tag{9}$$

and using the simplification $U = 0$ in (7), then the matter current is geodesic ($\dot{u}_a = u_{a;b}u^b = 0$), without shear ($\sigma_{ab} = u_{(a;b)} + \dot{u}_{(a}u_{b)} - \frac{1}{3}\theta h_{ab} = 0$) and without expansion ($\theta = u^a_{;a} = 0$); this implies that $A_{,r} = 0 = A_{,z}$. Moreover the dust is without rotation, $omega_{ab} = u_{[a;b]} + \dot{u}_{[a}u_{b]} = 0$.

If we take $A = 0, U = 0$ then

$$\left\{\begin{array}{l} g_{11} = g_{22} = e^{2V} = h_{11} = h_{22}, \\ g_{33} = W^2 = h_{33}, \quad g_{00} = \xi_0 = -1 = F, \\ g_{03} = \xi_3 = 0 = g_{13} = g_{23} = g_{10} = g_{20} \end{array}\right. \tag{10}$$

and the field equations (3) will have the following simplified form:

$$\left\{\begin{array}{l} \frac{1}{2}\Big(\frac{\partial^2 W}{\partial r^2} + \frac{\partial^2 W}{\partial z^2}\Big) = -\Lambda W e^{2V} \\ (\mu - 2\Lambda)e^{2V} = 0 \\ \frac{1}{2}\Big(\frac{\partial^2 W}{\partial r^2} + \frac{\partial^2 W}{\partial z^2}\Big) = \frac{\partial W}{\partial r}\frac{\partial V}{\partial r} - \frac{\partial W}{\partial z}\frac{\partial V}{\partial z} \\ \frac{\partial W}{\partial r}\frac{\partial V}{\partial z} + \frac{\partial W}{\partial z}\frac{\partial V}{\partial r} = 0 \\ \frac{\partial^2 V}{\partial r^2} + \frac{\partial^2 V}{\partial z^2} = -\Lambda e^{2V} \end{array}\right. \tag{11}$$

---

[2] The function W is defined invariantly as $W^2 \equiv -2\xi_{[a}\eta_{b]}\xi^a\eta^b$.

where W and V are functions of $x^1 = r$ and $x^2 = z$.

Finally, taking into account the third symmetry ($\zeta = \partial_z$ is the spacelike Killing vector field parallel with the rotation axis z) as a special case of the stationary axisymmetric (with ξ and η) solution (11) we will obtain:

$$\mathrm{d}s^2 = e^{2V(r)}(\mathrm{d}r^2 + \mathrm{d}z^2) + W^2(r)\mathrm{d}\varphi^2 - \mathrm{d}t^2 \tag{12}$$

where $W(r)$ and $V(r)$ will be determined from (11).

2 The solution

Because of the third symmetry W and V depend only on r and the field equations are reduced to:

$$\begin{cases} W'' = -2\Lambda W e^{2V} \\ W'' = 2W'V' \\ V'' + \Lambda e^{2V} = 0 \end{cases} \tag{13}$$

where $\frac{\partial}{\partial r} ='$ and $\mu = 2\Lambda = \text{const.}$.

The last equation in the (13) system is a differential equation of second order which implies:

$$r = \int \frac{1}{\sqrt{\gamma - \Lambda e^{2V}}}\,\mathrm{d}V + \alpha \tag{14}$$

where γ and α are constants.

Integrating in (14) we find:

$$r = \frac{1}{2\lambda}\ln\left|\frac{\sqrt{\gamma - \Lambda e^{2V}} - \lambda}{\sqrt{\gamma - \Lambda e^{2V}} + \lambda}\right| + \nu \tag{15}$$

Finally we obtain:

$$V(r) = \frac{1}{2}\ln\Big(\gamma - \lambda^2\Big(\frac{1 - e^{2\lambda(r-\nu)}}{1 + e^{2\lambda(r-\nu)}}\Big)^2\Big) - \ln\sqrt{\Lambda} \tag{16}$$

The first two equations in (13) reduce to

$$\frac{W'}{W} = -\Lambda\frac{e^{2V}}{V'} \tag{17}$$

which becomes:

$$\ln W = \lambda\int\frac{1 + e^{2\lambda(r-\nu)}}{1 - e^{2\lambda(r-\nu)}}\,\mathrm{d}r - \frac{\gamma}{\lambda^2}\int\frac{1 - e^{2\lambda(r-\nu)}}{1 + e^{2\lambda(r-\nu)}}\,\mathrm{d}r \tag{18}$$

Integrating both integrals we are lead to:

$$W(r) = \frac{1-e^{2\lambda(r-\nu)}}{\left(1-e^{2\lambda(r-\nu)}\right)^{\frac{\gamma}{\lambda^3}}}\, e^{2\lambda(r-\nu)(\frac{1}{2}-\frac{\gamma}{2\lambda^3})} \tag{19}$$

where γ, α, λ and ν are constants.

The resulting solution is static cylindrically symmetric and conformally flat and the matter consists in a cosmological and dust term with $\mu = 2\Lambda =$ const. positive.

Remark: Further asking the regularity condition on the axis of rotation to be satisfied, i.e.:

$$\lim_{r\to 0} \frac{\frac{1-e^{2\lambda(r-\nu)}}{\left(1-e^{2\lambda(r-\nu)}\right)^{\frac{\gamma}{\lambda^3}}}\, e^{2\lambda(r-\nu)(\frac{1}{2}-\frac{\gamma}{2\lambda^3})}}{r\sqrt{\lambda^2\left(\frac{1+e^{2\lambda(r-\nu)}}{1-e^{2\lambda(r-\nu)}}\right)^2-\gamma^2}} = 1 \tag{20}$$

we find out the regular Einstein's static universe (see [3]) solution in cylindrically coordinates (with $\mu = 2\Lambda = \frac{1}{2K^2} =$ const.).

References

[1] **Huesler, M.**, *Stationary Black Holes: uniqueness and beyond*, Living Reviews, (1998)

[2] **Kramer, D., Stephani, H., Herlt, E., MacCallum, M., Schmutzer, E.**, *Exact solutions of Einstein's field equations*, Cambridge Univ. Press, (1980)

[3] **Misner, C.W., Thorne, K.S., Wheeler, J.A.**, *Gravitation*, W.H. Freeman and Company, San Francisco (1973)